

The Full-sky Astrometric Mapping Explorer¹

Scott D. Horner^{2,3}, Marvin E. Germain⁴, Thomas P. Greene⁵, Fred H. Harris⁴, Mark S. Johnson⁶, Kenneth J. Johnston³, David G. Monet⁴, Marc A. Murison³, James D. Phillips⁷, Robert D. Reasenberg⁷, P. Kenneth Seidelmann³, Sean E. Urban³, and Richard H. Vassar⁸

FAME is designed to perform an all-sky, astrometric survey with unprecedented accuracy. It will create a rigid astrometric catalog of 4×10^7 stars with $5 < m_V < 15$. For bright stars, $5 < m_V < 9$, FAME will determine positions and parallaxes accurate to $< 50 \mu\text{as}$, with proper motion errors $< 50 \mu\text{as}/\text{year}$. For fainter stars, $9 < m_V < 15$, FAME will determine positions and parallaxes accurate to $< 500 \mu\text{as}$, with proper motion errors $< 500 \mu\text{as}/\text{year}$. It will also collect photometric data on these 4×10^7 stars in four Sloan DSS colors. NASA selected the Full-sky Astrometric Mapping Explorer (FAME) to be one of five MIDEX missions funded for a concept study. The Phase A Concept Study Report was submitted to NASA on 18 June 1999. In September 1999, NASA will select two of these five missions for flight as MIDEX-3 (CY2003 launch) and MIDEX-4 (CY2004 launch) in its Explorer program.

The greatest strength of FAME will be the large number of stars observed, enabling diverse studies of stellar and galactic evolution. FAME will provide a meaningful statistical sample of stars for studies of the frequency of stellar companions with $m > M_{\text{jupiter}}$. This will provide accurate statistics on the frequency of occurrence of multiple star systems as a function of spectral type, as well as the frequency of occurrence of giant planets. It will explore the possible transition region between giant planets and brown dwarfs, improving our understanding of star system formation. By determining accurate parallaxes for 4×10^7 stars, FAME will calibrate the absolute luminosities of solar neighborhood stars. This will determine the distances and ages of open and globular clusters, refine stellar structure models, and calibrate the distances to Cepheid and RR Lyrae stars. FAME will also monitor $\sim 40,000$ solar-like stars for photometric variations at the 0.1% level to search for evidence of magnetic activity cycles analogous to the 11-year solar activity cycle.

FAME will produce a major catalog of fundamental stellar properties that will enable advances across numerous branches of astrophysics. It will open up a new dimension in stellar astrophysics and map our quadrant of the Milky Way galaxy.

¹<http://www.usno.navy.mil/fame>

²shorner@usno.navy.mil

³U.S. Naval Observatory, 3450 Massachusetts Ave. NW, Washington, DC 20392-5420

⁴U.S. Naval Observatory, PO Box 1149, Flagstaff, AZ 86002-1149

⁵NASA Ames Research Center, M.S. 245-6, Moffett Field, CA 94035-1000

⁶Naval Research Laboratory, Code 8100, 4555 Overlook Ave. SW, Washington, DC 20375-5000

⁷Smithsonian Astrophysical Observatory, 60 Garden St., Cambridge, MA 02138

⁸Lockheed Martin Missiles and Space Advanced Technology Center, 3251 Hanover St., Palo Alto, CA 94304-1191




Figure 1. FAME will map our quadrant of the galaxy out to 2kpc from the Sun providing the information needed to calibrate the standard candles that define the extragalactic distance scale, calibrate the absolute luminosities of stars of all spectral types for studies of stellar structure and evolution, and detect orbital motions caused by brown dwarfs and giant planets.




Figure 2. Comparison of the astrometric capabilities of FAME to SIM and Hipparcos. The lines indicate where achieved accuracy will be 10% parallax error or better. Known Cepheids are indicated by circles and RR Lyraes with 'x's. Distances to clusters and OB associations are indicated at the right of the figure.